

# NOTES

Edited by William Adkins

---

## On Homeomorphism Groups and the Compact-Open Topology

---

Jan J. Dijkstra

---

If  $X$  is a topological space, then we let  $\mathcal{H}(X)$  denote the group of autohomeomorphisms of  $X$  equipped with the compact-open topology. For subsets  $A$  and  $B$  of  $X$  we define  $[A, B] = \{h \in \mathcal{H}(X) : h(A) \subset B\}$ , and we recall that the topology on  $\mathcal{H}(X)$  is generated by the subbasis  $\mathcal{S}_X = \{[K, O] : K \text{ compact, } O \text{ open in } X\}$ . If  $X$  is a compact Hausdorff space, then  $\mathcal{H}(X)$  is a topological group, that is, composition and taking inverses are continuous operations (see Arens [2]). It is well known that even for locally compact separable metric spaces the inverse operation on  $\mathcal{H}(X)$  may not be continuous, thus  $\mathcal{H}(X)$  is, in general, not a topological group (see the example to follow). However, it is a classic theorem of Arens [2] that if a Hausdorff space  $X$  is noncompact, locally compact, and locally connected, then  $\mathcal{H}(X)$  is a topological group because the compact-open topology coincides with the topology that  $\mathcal{H}(X)$  inherits from  $\mathcal{H}(\alpha X)$ , where  $\alpha X$  is the Alexandroff one-point compactification of  $X$ . We improve on this result as follows. (Recall that a *continuum* is a compact connected space. If  $A$  is a subset of  $X$ , then  $\text{int } A$  and  $\partial A$  denote the interior and boundary of  $A$  in  $X$ , respectively.)

**Theorem.** *Let  $X$  be a noncompact Hausdorff space. If every point in  $X$  has a neighbourhood that is a continuum, then the compact-open topology on  $\mathcal{H}(X)$  coincides with the group topology that  $\mathcal{H}(X)$  inherits from  $\mathcal{H}(\alpha X)$ .*

*Proof.* The topology that  $\mathcal{H}(X)$  inherits from  $\mathcal{H}(\alpha X)$  is generated by the subbasis

$$\mathcal{S}_X \cup \{[F, X \setminus K] : F \text{ closed, } K \text{ compact in } X\}$$

(see Arens [2], who calls this the *g-topology* on  $\mathcal{H}(X)$ ). Let  $F$  be closed in  $X$ , and let  $O$  be the complement of a compact subset  $K$  of  $X$ . We show that  $[F, O]$  is open in the compact-open topology. Let  $f$  be an arbitrary element of  $[F, O]$ , and consider the compactum  $f^{-1}(K)$ . Select for each  $x$  in  $f^{-1}(K)$  a neighbourhood  $C_x$  of  $x$  in  $X$  that is a continuum. By compactness we can find a finite subset  $A$  of  $f^{-1}(K)$  such that

$$f^{-1}(K) \subset \bigcup_{a \in A} \text{int } C_a.$$

Define the compactum  $C = \bigcup_{a \in A} C_a$  and construct in the same way a compact neighbourhood  $C'$  of  $C$ . Consider the obviously open subset

$$U = [C' \cap F, O] \cap [\partial C', f(X \setminus C)] \cap \bigcap_{a \in A} [\{a\}, f(\text{int } C_a)]$$

of  $\mathcal{H}(X)$  in the compact-open topology. It is clear that  $f$  belongs to  $U$ , and now we need verify only that  $U$  is a subset of  $[F, O]$ .

Assume to the contrary that  $h$  lies in  $U \setminus [F, O]$ . Thus there is an  $x$  in  $F$  with  $h(x)$  in  $K = X \setminus O$ . Since  $h \in U \subset [C' \cap F, O]$ , we see that  $x$  cannot lie in  $C'$ . Note that

$$x \in h^{-1}(K) \subset h^{-1}(f(C)),$$

hence  $x$  lies in  $h^{-1}(f(C_a))$  for some  $a$  in  $A$ . Since

$$h \in U \subset [\{a\}, f(\text{int } C_a)],$$

we see that  $a$  lies in  $h^{-1}(f(C_a))$ . Observe that  $h^{-1}(f(C_a))$  is a continuum that connects the point  $a$  inside  $C'$  with  $x$  outside  $C'$ , hence there is a  $y$  in  $h^{-1}(f(C_a)) \cap \partial C'$ . Since  $U \subset [\partial C', f(X \setminus C)]$ , we infer that  $h(y) \notin f(C)$ , which contradicts the fact that  $y$  is a member of  $h^{-1}(f(C_a))$ . The proof is complete. ■

**Example.** For the sake of completeness we include an example of a locally compact metric space  $X$  such that the inverse operation on  $\mathcal{H}(X)$  is not continuous. Let  $C$  be a Cantor set, and let  $a$  and  $b$  be distinct points of  $C$ . We consider the space  $\mathcal{H}(X)$  with the compact-open topology, where  $X = C \setminus \{a\}$ . Select for  $a$  and  $b$  neighbourhood bases  $\{U_n : n \in \mathbb{N}\}$  and  $\{V_n : n \in \mathbb{N}\}$ , respectively, such that  $U_1 \cap V_1 = \emptyset$ ,  $U_n$  and  $V_n$  are both closed and open, and  $U_{n+1} \subsetneq U_n$  and  $V_{n+1} \subsetneq V_n$  for each  $n$ . For instance, if we put

$$C = \left\{ \sum_{n=1}^{\infty} \varepsilon_n 3^{-n} : \varepsilon_n \in \{0, 2\} \text{ for each } n \in \mathbb{N} \right\},$$

that is,  $C$  is the “middle third” Cantor set in the real line, then we can simply take  $a = 0$ ,  $b = 1$ ,  $U_n = C \cap [0, 3^{-n}]$ , and  $V_n = C \cap [1 - 3^{-n}, 1]$ .

Noting that nonempty closed and open subsets of  $C$  are also Cantor sets, we can find for each  $n$  in  $\mathbb{N}$  an  $h_n$  in  $\mathcal{H}(C)$  that satisfies the following conditions:

$$\begin{aligned} h_n(x) &= x & (x \in C \setminus (U_n \cup V_n)), \\ h_n(a) &= a, \\ h_n(U_{n+1}) &= U_n, \\ h_n(U_n \setminus U_{n+1}) &= V_{n+1}, \\ h_n(V_n) &= V_n \setminus V_{n+1}. \end{aligned}$$

If  $C$  is the middle third set, then the map  $h_n$  can be realized easily via a piecewise linear transformation of the interval  $[0, 1]$ .

Since  $h_n(a) = a$ , we see that the restriction  $h_n \upharpoonright X$  is a member of  $\mathcal{H}(X)$  for each  $n$ . Let the element  $[K, O]$  of  $\mathcal{S}_X$  be a neighbourhood of the identity element  $e$  of  $\mathcal{H}(X)$ . Thus  $K$  is a compact subset of  $O$  and  $X$ , which enables us to find an  $M$  such that  $U_M$  and  $K$  are disjoint. If  $b$  is not a member of  $K$ , then there is an  $N$  such that  $N \geq M$  and  $V_N \cap K = \emptyset$ , so  $h_n \upharpoonright K$  is an identity map and  $h_n \upharpoonright X$  is a member of  $[K, O]$  for each  $n$  satisfying  $n \geq N$ . If  $b$  is a point in  $K$ , then there is an  $N$  such that  $N \geq M$  and  $V_N$  is contained in  $O$ . Consequently, whenever  $n \geq N$  we have

$$h_n(K \setminus V_n) = K \setminus V_n \subset O, \quad h_n(K \cap V_n) \subset V_n \subset V_N \subset O,$$

whence  $h_n \upharpoonright X$  is in  $[K, O]$ . We have shown that  $\lim_{n \rightarrow \infty} h_n \upharpoonright X = e$  in  $\mathcal{H}(X)$ . On the other hand,  $h_n^{-1}(b)$  lies in  $U_n$  for each  $n$ , so certainly  $\lim_{n \rightarrow \infty} (h_n \upharpoonright X)^{-1} \neq e$ .

**Remarks.** One of the more interesting (and challenging) problems in infinite-dimensional topology is the topological classification of  $\mathcal{H}(M)$ , where  $M$  is a manifold. Let  $\mathbb{I}$  denote the interval  $[0, 1]$ , and let  $\mathcal{H}_\partial(\mathbb{I}^n)$  stand for the subgroup of  $\mathcal{H}(\mathbb{I}^n)$  consisting of homeomorphisms that fix the boundary of the  $n$ -cube  $\mathbb{I}^n$ . Anderson [1] proved that  $\mathcal{H}_\partial(\mathbb{I})$  is homeomorphic to the separable Hilbert space  $\ell^2$  (see [3, Proposition VI.8.1] or [11]). It was shown by Luke and Mason [12] that  $\mathcal{H}_\partial(\mathbb{I}^2)$  is an absolute retract, that is, the space is homeomorphic to a retract of  $\ell^2$ . This result in combination with the fact that  $\mathcal{H}_\partial(\mathbb{I}^2)$  is a topological group guarantees that the space is homeomorphic to  $\ell^2$  (apply, for instance, Dobrowolski and Toruńczyk [7]). If  $n \geq 3$  it is wide open whether  $\mathcal{H}_\partial(\mathbb{I}^n)$  is an absolute retract. For the Hilbert cube  $Q$ , that is, for  $n = \infty$ , the corresponding problem was solved by Ferry [10] and Toruńczyk [13]. They proved that  $\mathcal{H}(Q)$  is homeomorphic to  $\ell^2$  (observe that  $Q$  has no boundary).

If  $D$  is a countable dense subset of  $M$ , then we let  $\mathcal{H}(M, D)$  signify the group  $\{h \in \mathcal{H}(M) : h(D) = D\}$ . The topological classification problem for  $\mathcal{H}(M, D)$  was recently solved completely in joint work of the author and Jan van Mill [4], [5], [6]. If  $M$  is a one-dimensional topological manifold or a Cantor set, then  $\mathcal{H}(M, D)$  is homeomorphic to  $\mathbb{Q}^\infty$ , the countable power of the space of rational numbers. If  $M$  is a topological manifold of dimension at least two, a Hilbert cube manifold, or a manifold modelled on a universal Menger continuum, then  $\mathcal{H}(M, D)$  is homeomorphic to the famed Erdős space [9], which consists of the vectors in  $\ell^2$  whose coordinates are all rational. (Menger continua are higher dimensional analogues of the Cantor set; see, for instance, [8, sec. 1.11].)

## REFERENCES

1. R. D. Anderson, Spaces of homeomorphisms of finite graphs (unpublished manuscript).
2. R. Arens, Topologies for homeomorphism groups, *Amer. J. Math.* **68** (1946) 593–610.
3. C. Bessaga and A. Pełczyński, *Selected Topics in Infinite-Dimensional Topology*, Polish Scientific Publishers, Warsaw, 1975.
4. J. J. Dijkstra and J. van Mill, Homeomorphism groups of manifolds and Erdős space, *Electron. Res. Announc. Amer. Math. Soc.* **10** (2004) 29–38.
5. ———, On the group of homeomorphisms of the real line that map the pseudoboundary onto itself, *Canad. J. Math.* (to appear).
6. ———, Erdős space and homeomorphism groups of manifolds (2004 preprint).
7. T. Dobrowolski and H. Toruńczyk, Separable complete ANRs admitting a group structure are Hilbert manifolds, *Topology Appl.* **12** (1981) 229–235.
8. R. Engelking, *Dimension Theory*, North-Holland, Amsterdam, 1978.
9. P. Erdős, The dimension of the rational points in Hilbert space, *Ann. of Math.* **41** (1940) 734–736.
10. S. Ferry, The homeomorphism group of a compact Hilbert cube manifold is an ANR, *Ann. of Math.* **106** (1977) 101–119.
11. J. E. Keesling, Using flows to construct Hilbert space factors of function spaces, *Trans. Amer. Math. Soc.* **161** (1971) 1–24.
12. R. Luke and W. K. Mason, The space of homeomorphisms of a compact two-manifold is an absolute neighborhood retract, *Trans. Amer. Math. Soc.* **164** (1972) 275–285.
13. H. Toruńczyk, Homeomorphism groups of compact Hilbert cube manifolds which are manifolds, *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.* **25** (1977) 401–408.

*Faculteit der Exacte Wetenschappen/Afdeling Wiskunde, Vrije Universiteit, De Boelelaan 1081a, 1081 HV Amsterdam, The Netherlands*  
*dijkstra@cs.vu.nl*