

advanced logic

2019 02 11

lecture 3

so far and alternatives

formulas: quite a few alternative approaches possible and used

for example: negation only on atoms and further defined inductively

valuation: alternative approach could be (not used)

$$\sigma : W \rightarrow \mathcal{P}(\text{Var})$$

semantics: Kripke models

alternative approaches possible and used

semantics in terms of sets of worlds, game semantics

overview

- preservation of truth and validity
- characterizations of frame properties
- bisimulations

overview

- preservation of truth and validity
- characterizations of frame properties
- bisimulations

prop1: schemes preservation under substitution

sometimes we consider a formula as a scheme:

$$p \rightarrow q \rightarrow p$$

$$(r \wedge s) \rightarrow \perp \rightarrow (r \wedge s)$$

let $\sigma : Var \rightarrow Form$ be a substitution and $v : Var \rightarrow \{0, 1\}$ a valuation

if $\llbracket \phi \rrbracket_v = 1$ then not necessarily $\llbracket \phi^\sigma \rrbracket_v = 1$

if $\llbracket \phi \rrbracket = 1$ (or: $\models \phi$) then $\models \phi^\sigma$

definition: substitution for propositional variables

a mapping $\sigma : \text{Var} \rightarrow \text{Form}$ induces a mapping $(\cdot)^\sigma : \text{Form} \rightarrow \text{Form}$:

$$p^\sigma = \sigma(p)$$

$$\perp^\sigma = \perp$$

$$\top^\sigma = \top$$

$$(X\phi)^\sigma = X(\phi^\sigma) \quad \text{for } X \in \{\neg, \Box, \Diamond\}$$

$$(\phi * \psi)^\sigma = (\phi^\sigma) * (\psi^\sigma) \quad \text{for } * \in \{\vee, \wedge, \rightarrow\}$$

modal prop1: what validity is preserved under substitution?

let $\sigma : Var \rightarrow Form$ be a substitution

if $((W, R), V), w \models \phi$ then not necessarily $((W, R), V), w \models \phi^\sigma$

if $((W, R), V) \models \phi$ then not necessarily $((W, R), V) \models \phi^\sigma$

if $(W, R) \models \phi$ then $(W, R) \models \phi^\sigma$?

if so, then also: if $\models \phi$ then $\models \phi^\sigma$

validity in a frame is preserved under substitution

we will see:

if $\mathcal{F} \models \phi$ then $\mathcal{F} \models \phi^\sigma$ for any substitution σ

how do we prove this?

validity in a frame is preserved under substitution

we will see:

if $\mathcal{F} \models \phi$ then $\mathcal{F} \models \phi^\sigma$ for any substitution σ

how do we prove this?

assume $\mathcal{F} \models \phi$ with $\mathcal{F} = (W, R)$ and let σ be a substitution

validity in a frame is preserved under substitution

we will see:

if $\mathcal{F} \models \phi$ then $\mathcal{F} \models \phi^\sigma$ for any substitution σ

how do we prove this?

assume $\mathcal{F} \models \phi$ with $\mathcal{F} = (W, R)$ and let σ be a substitution

let V be a valuation and let $w \in W$

we aim to show $\mathcal{F}, V, w \models \phi^\sigma$

validity in a frame is preserved under substitution

we will see:

if $\mathcal{F} \models \phi$ then $\mathcal{F} \models \phi^\sigma$ for any substitution σ

how do we prove this?

assume $\mathcal{F} \models \phi$ with $\mathcal{F} = (W, R)$ and let σ be a substitution

let V be a valuation and let $w \in W$

we aim to show $\mathcal{F}, V, w \models \phi^\sigma$

we would like to use $\mathcal{F}, V^\sigma, w \models \phi$

with V^σ mapping p to the set of worlds in which $\sigma(p)$ is true

use 'alternative semantics'

recall: alternative semantics

Let $\mathcal{M} = (W, R, V)$ be a model.

We define $\llbracket \phi \rrbracket_{\mathcal{M}} \subseteq W$, the *interpretation* of a formula ϕ in the model \mathcal{M} , inductively by

$$\llbracket p \rrbracket_{\mathcal{M}} = V(p) \quad (p \in \text{Var})$$

$$\llbracket \perp \rrbracket_{\mathcal{M}} = \emptyset$$

$$\llbracket \top \rrbracket_{\mathcal{M}} = W$$

$$\llbracket \neg \phi \rrbracket_{\mathcal{M}} = W \setminus \llbracket \phi \rrbracket_{\mathcal{M}}$$

$$\llbracket \phi \vee \psi \rrbracket_{\mathcal{M}} = \llbracket \phi \rrbracket_{\mathcal{M}} \cup \llbracket \psi \rrbracket_{\mathcal{M}}$$

$$\llbracket \phi \wedge \psi \rrbracket_{\mathcal{M}} = \llbracket \phi \rrbracket_{\mathcal{M}} \cap \llbracket \psi \rrbracket_{\mathcal{M}}$$

$$\llbracket \phi \rightarrow \psi \rrbracket_{\mathcal{M}} = \neg \llbracket \phi \rrbracket_{\mathcal{M}} \cup \llbracket \psi \rrbracket_{\mathcal{M}}$$

$$\llbracket \diamond \phi \rrbracket_{\mathcal{M}} = \{w \in W \mid \exists v R w v \wedge v \in \llbracket \phi \rrbracket_{\mathcal{M}}\}$$

$$\llbracket \square \phi \rrbracket_{\mathcal{M}} = \{w \in W \mid \forall v R w v \Rightarrow v \in \llbracket \phi \rrbracket_{\mathcal{M}}\}$$

(for $X \subseteq W$ we write $\neg X$ to denote the *complement* of X , i.e.,
 $\neg X = W \setminus X$)

lemma: intuition about interpretation is correct

the interpretation $\llbracket \phi \rrbracket_{\mathcal{M}}$ of formula ϕ in model $\mathcal{M} = (W, R, V)$

is the set of worlds in which ϕ is true:

Lemma: $\mathcal{M}, w \models \phi$ if and only if $w \in \llbracket \phi \rrbracket_{\mathcal{M}}$

consequence of the lemma:

$\mathcal{M} \models \phi$ if and only if $\llbracket \phi \rrbracket_{\mathcal{M}} = W$

evaluating substitution instances

let $\mathcal{M} = (W, R, V)$ be a model, and let $\sigma : Var \rightarrow Form$ be a substitution

definition: a substitution σ applied to a valuation:

$$V^\sigma(p) = \llbracket \sigma(p) \rrbracket_{\mathcal{M}}$$

Lemma:

$$\llbracket \phi^\sigma \rrbracket_{(W,R,V)} = \llbracket \phi \rrbracket_{(W,R,V^\sigma)}$$

theorem: validity is closed under substitution

if $\mathcal{F} \models \phi$ then $\mathcal{F} \models \phi^\sigma$ for any substitution σ

assume $\mathcal{F} \models \phi$ with $\mathcal{F} = (W, R)$ and let σ be a substitution

we have to show $\mathcal{F} \models \phi^\sigma$

let V be a valuation and let $w \in W$

we have $\mathcal{F}, V^\sigma, w \models \phi$

hence $w \in \llbracket \phi \rrbracket_{\mathcal{F}, V^\sigma}$

hence $w \in \llbracket \phi^\sigma \rrbracket_{\mathcal{F}, V}$

hence $\mathcal{F}, V, w \models \phi^\sigma$

hence $\mathcal{F} \models \phi^\sigma$

consequences of this theorem

take a propositional tautology, apply a substitution, then we have a modal tautology

how can we use this?

for example: consider the substitution δ with $\delta(p) = \neg p$

$\mathcal{F} \models p \rightarrow \diamond p$ if and only if $\mathcal{F} \models \Box p \rightarrow p$

$\mathcal{F} \models \diamond p \rightarrow \Box \diamond p$ if and only if $\mathcal{F} \models \diamond \Box p \rightarrow \Box p$

preservation of truth and validity

local truth is preserved by modus ponens:

if $\mathcal{M}, w \models \phi \rightarrow \psi$ and $\mathcal{M}, w \models \phi$ then $\mathcal{M}, w \models \psi$

global truth is preserved by modus ponens and by necessitation:

if $\mathcal{M} \models \phi$ then $\mathcal{M} \models \Box\phi$

frame validity is preserved by modus ponens, necessitation, and substitution:

if $\mathcal{F} \models \phi$ then $\mathcal{F} \models \phi^\sigma$

the modal tautologies are exactly defined by

extension:

a tautology for first-order propositional logic is a modal tautology

modal distribution:

$$\vDash \Box(p \rightarrow q) \rightarrow \Box p \rightarrow \Box q$$

modus ponens:

if $\vDash \phi \rightarrow \psi$ and $\vDash \phi$ then $\vDash \psi$

necessitation:

if $\vDash \phi$ then $\vDash \Box \phi$

substitution:

if $\vDash \phi$ then $\vDash \phi^\sigma$

example frame characterization

if $\mathcal{F} = (W, R)$ is symmetric then $\mathcal{F} \models \diamond\Box p \rightarrow p$

example frame characterization

if $\mathcal{F} = (W, R)$ is symmetric then $\mathcal{F} \models \diamond\Box p \rightarrow p$

let $\mathcal{F} = (W, R)$ be a symmetric frame

example frame characterization

if $\mathcal{F} = (W, R)$ is symmetric then $\mathcal{F} \models \diamond\Box p \rightarrow p$

let $\mathcal{F} = (W, R)$ be a symmetric frame

let $\mathcal{M} = (\mathcal{F}, V)$ be a model based on \mathcal{F} , and let $w \in W$

example frame characterization

if $\mathcal{F} = (W, R)$ is symmetric then $\mathcal{F} \models \diamond\Box p \rightarrow p$

let $\mathcal{F} = (W, R)$ be a symmetric frame

let $\mathcal{M} = (\mathcal{F}, V)$ be a model based on \mathcal{F} , and let $w \in W$

assume $\mathcal{M}, w \models \diamond\Box p$; we have to show $\mathcal{M}, w \models p$

example frame characterization

if $\mathcal{F} = (W, R)$ is symmetric then $\mathcal{F} \models \Diamond \Box p \rightarrow p$

let $\mathcal{F} = (W, R)$ be a symmetric frame

let $\mathcal{M} = (\mathcal{F}, V)$ be a model based on \mathcal{F} , and let $w \in W$

assume $\mathcal{M}, w \models \Diamond \Box p$; we have to show $\mathcal{M}, w \models p$

because $w \models \Diamond \Box p$ there is a $v \in W$ with Rwv and $v \models \Box p$

example frame characterization

if $\mathcal{F} = (W, R)$ is symmetric then $\mathcal{F} \models \Diamond \Box p \rightarrow p$

let $\mathcal{F} = (W, R)$ be a symmetric frame

let $\mathcal{M} = (\mathcal{F}, V)$ be a model based on \mathcal{F} , and let $w \in W$

assume $\mathcal{M}, w \models \Diamond \Box p$; we have to show $\mathcal{M}, w \models p$

because $w \models \Diamond \Box p$ there is a $v \in W$ with Rwv and $v \models \Box p$

because \mathcal{F} is symmetric we have Rvw

example frame characterization

if $\mathcal{F} = (W, R)$ is symmetric then $\mathcal{F} \models \diamond\Box p \rightarrow p$

let $\mathcal{F} = (W, R)$ be a symmetric frame

let $\mathcal{M} = (\mathcal{F}, V)$ be a model based on \mathcal{F} , and let $w \in W$

assume $\mathcal{M}, w \models \diamond\Box p$; we have to show $\mathcal{M}, w \models p$

because $w \models \diamond\Box p$ there is a $v \in W$ with Rwv and $v \models \Box p$

because \mathcal{F} is symmetric we have Rvw

because $v \models \Box p$ we have $w \models p$

example frame characterization

if $\mathcal{F} = (W, R)$ is symmetric then $\mathcal{F} \models \Diamond \Box p \rightarrow p$

let $\mathcal{F} = (W, R)$ be a symmetric frame

let $\mathcal{M} = (\mathcal{F}, V)$ be a model based on \mathcal{F} , and let $w \in W$

assume $\mathcal{M}, w \models \Diamond \Box p$; we have to show $\mathcal{M}, w \models p$

because $w \models \Diamond \Box p$ there is a $v \in W$ with Rwv and $v \models \Box p$

because \mathcal{F} is symmetric we have Rvw

because $v \models \Box p$ we have $w \models p$

so $\mathcal{M}, w \models \Diamond \Box p \rightarrow p$

example frame characterization

if $\mathcal{F} = (W, R)$ is symmetric then $\mathcal{F} \models \Diamond \Box p \rightarrow p$

let $\mathcal{F} = (W, R)$ be a symmetric frame

let $\mathcal{M} = (\mathcal{F}, V)$ be a model based on \mathcal{F} , and let $w \in W$

assume $\mathcal{M}, w \models \Diamond \Box p$; we have to show $\mathcal{M}, w \models p$

because $w \models \Diamond \Box p$ there is a $v \in W$ with Rwv and $v \models \Box p$

because \mathcal{F} is symmetric we have Rvw

because $v \models \Box p$ we have $w \models p$

so $\mathcal{M}, w \models \Diamond \Box p \rightarrow p$

so $\mathcal{F} \models \Diamond \Box p \rightarrow p$

example frame characterization

if $\mathcal{F} \models \diamond\Box p \rightarrow p$ then \mathcal{F} is symmetric

example frame characterization

if $\mathcal{F} \models \diamond\Box p \rightarrow p$ then \mathcal{F} is symmetric

assume $\mathcal{F} = (W, R)$ is not symmetric; we will show $\mathcal{F} \not\models \diamond\Box p \rightarrow p$

example frame characterization

if $\mathcal{F} \models \Diamond\Box p \rightarrow p$ then \mathcal{F} is symmetric

assume $\mathcal{F} = (W, R)$ is not symmetric; we will show $\mathcal{F} \not\models \Diamond\Box p \rightarrow p$

because \mathcal{F} is not symmetric there are $a, b \in W$ such that Rab but not Rba

(note: a non-symmetric frame has at least two states)

example frame characterization

if $\mathcal{F} \models \Diamond\Box p \rightarrow p$ then \mathcal{F} is symmetric

assume $\mathcal{F} = (W, R)$ is not symmetric; we will show $\mathcal{F} \not\models \Diamond\Box p \rightarrow p$

because \mathcal{F} is not symmetric there are $a, b \in W$ such that Rab but not Rba

(note: a non-symmetric frame has at least two states)

define V such that $V(p) = \{x \in W \mid Rbx\}$ and let $\mathcal{M} = (\mathcal{F}, V)$

example frame characterization

if $\mathcal{F} \models \Diamond \Box p \rightarrow p$ then \mathcal{F} is symmetric

assume $\mathcal{F} = (W, R)$ is not symmetric; we will show $\mathcal{F} \not\models \Diamond \Box p \rightarrow p$

because \mathcal{F} is not symmetric there are $a, b \in W$ such that Rab but not Rba

(note: a non-symmetric frame has at least two states)

define V such that $V(p) = \{x \in W \mid Rbx\}$ and let $\mathcal{M} = (\mathcal{F}, V)$

then we have $\mathcal{M}, b \models \Box p$

example frame characterization

if $\mathcal{F} \models \Diamond \Box p \rightarrow p$ then \mathcal{F} is symmetric

assume $\mathcal{F} = (W, R)$ is not symmetric; we will show $\mathcal{F} \not\models \Diamond \Box p \rightarrow p$

because \mathcal{F} is not symmetric there are $a, b \in W$ such that Rab but not Rba

(note: a non-symmetric frame has at least two states)

define V such that $V(p) = \{x \in W \mid Rbx\}$ and let $\mathcal{M} = (\mathcal{F}, V)$

then we have $\mathcal{M}, b \models \Box p$

because Rab we have $\mathcal{M}, a \models \Diamond \Box p$

example frame characterization

if $\mathcal{F} \models \Diamond \Box p \rightarrow p$ then \mathcal{F} is symmetric

assume $\mathcal{F} = (W, R)$ is not symmetric; we will show $\mathcal{F} \not\models \Diamond \Box p \rightarrow p$

because \mathcal{F} is not symmetric there are $a, b \in W$ such that Rab but not Rba

(note: a non-symmetric frame has at least two states)

define V such that $V(p) = \{x \in W \mid Rbx\}$ and let $\mathcal{M} = (\mathcal{F}, V)$

then we have $\mathcal{M}, b \models \Box p$

because Rab we have $\mathcal{M}, a \models \Diamond \Box p$

because not Rba we have $a \notin V(p)$ so we have $\mathcal{M}, a \not\models p$

example frame characterization

if $\mathcal{F} \models \Diamond\Box p \rightarrow p$ then \mathcal{F} is symmetric

assume $\mathcal{F} = (W, R)$ is not symmetric; we will show $\mathcal{F} \not\models \Diamond\Box p \rightarrow p$

because \mathcal{F} is not symmetric there are $a, b \in W$ such that Rab but not Rba

(note: a non-symmetric frame has at least two states)

define V such that $V(p) = \{x \in W \mid Rbx\}$ and let $\mathcal{M} = (\mathcal{F}, V)$

then we have $\mathcal{M}, b \models \Box p$

because Rab we have $\mathcal{M}, a \models \Diamond\Box p$

because not Rba we have $a \notin V(p)$ so we have $\mathcal{M}, a \not\models p$

so $\mathcal{M}, a \not\models \Diamond\Box p \rightarrow p$

example frame characterization

if $\mathcal{F} \models \Diamond\Box p \rightarrow p$ then \mathcal{F} is symmetric

assume $\mathcal{F} = (W, R)$ is not symmetric; we will show $\mathcal{F} \not\models \Diamond\Box p \rightarrow p$

because \mathcal{F} is not symmetric there are $a, b \in W$ such that Rab but not Rba

(note: a non-symmetric frame has at least two states)

define V such that $V(p) = \{x \in W \mid Rbx\}$ and let $\mathcal{M} = (\mathcal{F}, V)$

then we have $\mathcal{M}, b \models \Box p$

because Rab we have $\mathcal{M}, a \models \Diamond\Box p$

because not Rba we have $a \notin V(p)$ so we have $\mathcal{M}, a \not\models p$

so $\mathcal{M}, a \not\models \Diamond\Box p \rightarrow p$

so $\mathcal{F} \not\models \Diamond\Box p \rightarrow p$

frame characterization

we have shown:

the formula $\Diamond\Box p \rightarrow p$ characterizes the frame property **symmetry**

in general:


the formula ϕ characterizes the frame property P means

\mathcal{F} has property P if and only if $\mathcal{F} \models \phi$

overview

- preservation of truth and validity
- characterizations of frame properties
- bisimulations

distinguishable and indistinguishable states



we can distinguish state 1 from state 3

we do not succeed in distinguishing state 2 from state 4

can we prove that it is not possible to distinguish state 2 from state 4?

bisimulation: definition

Let $\mathcal{M} = (W, R, V)$ and $\mathcal{M}' = (W', R', V')$ be models.

A non-empty relation $Z \subseteq W \times W'$ is a **bisimulation**,

notation $Z : \mathcal{M} \leftrightarrow \mathcal{M}'$,

if for all pairs $(w, w') \in Z$ we have the following:

- $w \in V(p)$ if and only if $w' \in V'(p)$
- if Rwv then for some $v' \in W'$ we have $R'w'v'$ and vZv'
- if $R'w'v'$ then for some $v \in W$ we have Rwv and vZv'

bisimulation: base



$$\mathcal{M} = (W, R, V)$$

$$\mathcal{M}' = (W', R', V')$$

if wZw' then for all $p \in \text{Var}$ we have $w \in V(p)$ if and only if $w' \in V'(p)$

bisimulation: base



$$\mathcal{M} = (W, R, V)$$

$$\mathcal{M}' = (W', R', V')$$

if wZw' then for all $p \in \text{Var}$ we have $w \in V(p)$ if and only if $w' \in V'(p)$

bisimulation: zig




$\mathcal{M} = (W, R, V)$

$\mathcal{M}' = (W', R', V')$


if wZw' and Rwv then there exists $v' \in W'$ such that $R'w'v'$ and vZv'

bisimulation: zig




if wZw' and Rwv then there exists $v' \in W'$ such that $R'w'v'$ and vZv'

bisimulation: zig




if wZw' and Rwv then there exists $v' \in W'$ such that $R'w'v'$ and vZv'

bisimulation: zig



if wZw' and Rwv then there exists $v' \in W'$ such that $R'w'v'$ and vZv'

bisimulation: zig



if wZw' and Rwv then there exists $v' \in W'$ such that $R'w'v'$ and vZv'

bisimulation: zag




$\mathcal{M} = (W, R, V)$

$\mathcal{M}' = (W', R', V')$


if wZw' and $R'w'v'$ then there exists $v \in W$ such that Rwv and vZv'

bisimulation: zag




if wZw' and $R'w'v'$ then there exists $v \in W$ such that Rwv and vZv'

bisimulation: zag




if wZw' and $R'w'v'$ then there exists $v \in W$ such that Rwv and vZv'

bisimulation: zag



if wZw' and $R'w'v'$ then there exists $v \in W$ such that Rwv and vZv'

bisimulation: zag



if wZw' and $R'w'v'$ then there exists $v \in W$ such that Rwv and vZv'

bisimilarity: definition

two models $\mathcal{M} = (W, R, V)$ and $\mathcal{M}' = (W', R', V')$ are bisimilar,

notation $\mathcal{M} \Leftrightarrow \mathcal{M}'$,


if there exists a bisimulation $Z \subseteq W \times W'$

two pointed models \mathcal{M}, w and \mathcal{M}', w' are bisimilar,

notation $\mathcal{M}, w \Leftrightarrow \mathcal{M}', w'$ or $w \Leftrightarrow w'$,


if there exists a bisimulation $Z : \mathcal{M} \Leftrightarrow \mathcal{M}'$ such that $(w, w') \in Z$

example bisimilar states



states 2 and 4 are bisimilar

example bisimilar states



states 2 and 4 are bisimilar

$$B_1 = \{(2, 4), (3, 3)\}$$

$$B_2 = \{(1, 1), (2, 4), (4, 2), (3, 3)\}$$

$$B_3 = \{(1, 1), (2, 2), (2, 4), (3, 3), (4, 2), (4, 4)\}$$

example bisimilar states

$$\mathcal{N} = (\mathbb{N}, S)$$

$$S = \{(n, n+1) \mid n \in \mathbb{N}\}$$

$$V(p) = \{2n \mid n \in \mathbb{N}\}$$



$$\mathcal{F} = (\{e, o\}, R)$$

$$R = \{(e, o), (o, e)\}$$


$$U(p) = \{e\}$$

the states 0 and e are bisimilar


non-bisimilarity: example




non-bisimilarity: example



non-bisimilarity: example



non-bisimilarity: example



non-bisimilarity: example

